



**Kerala Agricultural University, Vellanikkara, Thrissur**  
**Department of Agricultural Meteorology, College of Horticulture**

Email: cohagmet@kau.in & kauagmet@yahoo.co.in

**Agromet Advisory Service Bulletin**

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



**PALAKKAD DISTRICT**

**AAS 03/2019(5), Tuesday**

**15.01.2019**

**CURRENT SYNOPTIC SITUATION:** A trough in easterlies extending upto 0.9 km above mean sea level runs from Maldives area to southeast Arabian sea off Karnataka coast .

**WEATHER FORECAST FOR THE NEXT FIVE DAYS:**

Weather parameters	16.01.2019	17.01.2019	18.01.2019	19.01.2019	20.01.2019
Rainfall (mm)	0	0	0	0	0
Maximum temperature (°C )	31	31	31	31	31
Minimum temperature (°C )	22	22	22	22	22
Total cloud cover (octa)	6	3	4	5	5
Maximum Relative Humidity (%)	80	80	80	80	80
Minimum Relative Humidity (%)	35	35	35	35	35
Wind speed (kmph)	14	1.4	14	14	10
Wind direction (deg)	90	90	90	90	90

**No rainfall is forecasted for coming five days in the district by India Meteorological Department.**

**AGRO-METEOROLOGICAL ADVISORIES:**

Crop	Crop stage	Pest/ Disease	Advisories
Paddy	Milky stage	Rice bug	There is a chance of appearance of rice bug in paddy. To control them, spray 2 ml malathion per one litre of water. Pesticides application should be done either before 9 am or after 3 pm. Otherwise, the pesticide application will adversely affect the pollination.
Coconut	-	Red palm weevil	Coconut palms are seen affected by red palm weevil. Remove the affected parts from crown and apply one ml imidacloprid /one litre of water at spindle leaf area.
Vegetables (cowpea)	-	Fruit fly	There is a chance of fruit fly in vegetables. Remove and destroy affected and decayed fruits. Use special pheromone traps like cue lure (6 nos/Acre) for vegetable fruit fly along with Thulasi/fruit traps. If the attack is not reduced, spray mixture of 2 ml malathion and 10 g jaggery in one litre of water during flowering and fruit initiation stage.
Animal Husbandry	Mix 30g sodium bicarbonate and 1 teaspoon yeast in the feed daily to avoid digestive disorders associated with excessive acidity. (CAADECCS, KVASU, Mannuthy)		