




Agromet Advisory Service bulletin 20/18(40), Friday dt .18.05.2018
(Issued jointly by IAAS Vellanikkara)



Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara
E mail: kauagmet@yahoo.co.in, cohagmet@kau.in

THRISSUR

CURRENT SYNOPTIC SITUATION: A cyclonic circulation lies over Lakshadweep area and neighbourhood extending upto 3.1 km above mean sea level.

Heavy Rainfall Warning: Heavy rainfall is most likely to occur at one or two places in Kerala upto 20th May 2018.

INDIA METEOROLOGICAL DEPARTMENT
NWP MODELS BASED DISTRICT LEVEL WEATHER PREDICTION
ISSUED ON: 18-05-2018
VALID TILL 08:30 IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS	WEATHER FCST				
	DAY-1 19/05/2018	DAY-2 20/05/2018	DAY-3 21/05/2018	DAY-4 22/05/2018	DAY-5 23/05/2018
DISTRICT : THRISSUR					
Rainfall (mm)	8	9	8	10	14
Max Temperature (deg C)	33	33	33	33	33
Min Temperature (deg C)	22	22	22	22	22
Total cloud cover (octa)	7	7	7	7	7
Max Relative Humidity (%)	90	90	90	90	90
Min Relative Humidity (%)	70	70	70	70	70
Wind speed (kmph)	010	010	010	010	010
Wind direction (deg)	320	320	320	320	320

Agro Meteorological Advisories:

Crop	Crop stage / Pest/Disease	Advisories
Paddy(<i>puncha</i>)	Maturity stage	Paddy is in the harvesting stage.
Banana	Sigatoka	Due to the changing weather pattern, there is a chance of sigatoka disease in banana. As a precaution, Spray Pseudomonas 10-20 g per 1 liter of water. In case of severe attack spray 1 gm Bavistin by mixing it with gum in 1 litre of water on lower surface of leaf.
Pepper	Quick wilt	There is a chance of Quick wilt disease due to high relative humidity. Apply trichoderma or pseudomonas culture in soil or 1% Bordeaux mixture can be sprayed on the leaves and small branches.
Animal Husbandry	Adopt measures to eradicate external parasitic infestations in animals. (CAADECCS, KVASU, Mannuthy)	

Preparation of Trichoderma culture: Dry neem cake and powdered cow dung are mixed at 1:10 ratio and add 1 kg trichoderma with 100 kg of above mixture. Then sprinkle water and mix it thoroughly. Cover it with a perforated polythene sheet and keep it in shade for 5 days for multiplication. Again mix well and keep for 3 more days for further multiplication and incorporate into the soil for controlling Quick wilt.