



Kerala Agricultural University, Vellanikkara, Thrissur
Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in & kauagmet@yahoo.co.in

Agromet Advisory Service Bulletin

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



THRISSUR DISTRICT

AAS 28/2019(56), Friday

12.07.2019

SIGNIFICANT WEATHER FOR THE PRECEDING WEEK AT VELLANIKKARA:

The total rainfall received 159.6 mm. The maximum temperature ranges from 30.5°C to 32.6°C, minimum temperature ranges from 21.6°C to 23.8°C. Morning relative humidity ranges from 85 % to 98% and afternoon relative humidity ranges from 65% to 90%. Wind speed ranges from 0.7 to 3.4 km/h. Bright sunshine hours range from 0.5 to 6.1 hours. Evaporation ranges from 1.8 to 3.4 mm.

CURRENT SYNOPTIC SITUATION:

A feeble off shore trough runs from Karnataka coast to North Kerala coast. An upper air cyclonic circulation lies over 3.1 & 5.8km above mean sea level.

Heavy Rainfall Warning:

Day 5 : Heavy rainfall (7- 11 cm in 24 hours) is very likely to occur at isolated places in the district of Malappuram.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	13.07.2019	14.07.2019	15.07.2019	16.07.2019	17.07.2019
Rainfall (mm)	0	15	8	1	5
Maximum temperature (°C)	31	31	31	31	31
Minimum temperature (°C)	23	23	23	23	23
Maximum Relative Humidity (%)	98	98	98	98	98
Minimum Relative Humidity (%)	75	75	75	75	75
Wind speed (kmph)	6	6	6	6	8
Wind direction (deg)	270	270	290	290	290
Total cloud cover (octa)	5	8	7	6	7

There is a chance of reduction in rainfall compared to previous week in the district by IMD & KAU.

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Stage	Pest/ Disease	Advisories
Rice (Virippu)	Tillering stage	Leaf folder	Since cloudy weather is prevailed, there is a chance of Leaf folder attack in <i>Virippu</i> Paddy. It can be controlled by using trichogramma cards (2CC per one acre). Cut the cards in to small pieces and fix in different regions of the field. If the attack is severe, spray quinalphos @ 2ml per one liter of water.
Coconut		Bud rot	There is a chance of bud rot in rainy season. Keep three numbers each of perforated Mancozeb sachets (5 gm) around spindle leaves as a prophylactic measure against bud rot.
Banana	Bunch emergence stage	Sigatoka disease	Due to the pre monsoon showers, there is a chance of Sigatoka disease in banana. As a precaution, Spray Pseudomonas 20g/litre. In case of severe attack spray 1 gm Bavistin by mixing it with gum in 1 litre of water on lower surface of leaf.
Vegetables	Flowering/Fruiting stage	Downy mildew	There is a chance of downy mildew attack in vegetables due to the south west monsoon. Application of 0.2% curzate M-8 or 0.1% Equation pro or Trichoderma 20g/litre under the leaves in 15 days interval is effective.
Animal Husbandry			Don't allow the animals to water logged areas to avoid foot lesions. (CAADECCS, KVASU, Mannuthy)