



THRISSUR

Significant weather for the preceding week at Vellanikkara

Rainfall weekly total	0.0 mm
Maximum temperature	32.2-34.30 ⁰ C
Minimum temperature	19.4-22.3 ⁰ C
Max. RH	88-95%
Min. RH	46-71%
Wind speed	0.5-1.2 km/hr
Sun shine hour	5.8-9.2 hr
Evaporation	2.2-2.8 mm

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST
ISSUED ON: 12.12.2017
VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

CURRENT SYNOPTIC SITUATION: An upper air cyclonic circulation lies over Comorin area & neighbourhood and extends upto 0.9 km above mean sea level.

DISTRICT : THRISSUR

ENSEMBLE FCST

PARAMETERS	13.12.2017	14.12.2017	15.12.2017	16.12.2017	17.12.2017
Rainfall (mm)	2	0	0	0	0
Max Temperature (deg	31	31	31	31	31
Min Temperature (deg C)	21	21	21	21	21
Total cloud cover (octa)	5	4	4	4	4
Max Relative Humidity (%)	85	85	85	85	85
Min Relative Humidity (%)	75	75	75	75	75
Wind speed (kmph)	10	10	10	10	10
Wind direction (deg)	90	90	90	90	90

Agro Meteorological Advisories:

Crop	Crop stage / Pest/Disease	Advisories
Paddy (Jyothi, Kanchana)	Rice bug	Apply neem based pesticides (3ml per one liter of water). Pesticides application should be done either before 9 am or after 3 pm. Otherwise, the pesticide application will adversely affect the pollination.
Coconut	Bud rot	As a prophylactic measure, hung three perforated packets of Mancozeb sachet (5gm) around spindle leaves.
Banana	Sigatoka disease	Due to continuous rain, there is a chance of Sigatoka disease in banana. As a precaution, Spray Pseudomonas 20g/litre. In case of severe attack, spray 1 gm of Bavistin in 1 litre of water on lower surface of leaf (by mixing with gum)
Vegetables	Thrips and Mite	There is a chance for the attack of thrips and mites in vegetables. Apply 2% neem oil+garlic emulsion under the leaf surface at 10 days intervals. If the attack is severe apply spiromecifen 8 ml/10 L of water.
Animal Husbandry	If respiratory distress, fever or lameness are noticed in animals, consult a veterinary surgeon immediately. (CAADECS, KVASU, Mannuthy)	