





Agromet Advisory Bulletin for Ernakulam District (Valid from 20.01.2024 to 24.01.2024)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 03/2024, (6) Friday

Dated 19.01.2024

A. Medium range weather forecast of IMD for next five days

| Weather parameters | 20.01.2024 | 21.01.2024 | 22.01.2024 | 23.01.2024 | 24.01.2024 |
|-------------------------------|------------|------------|------------|------------|------------|
| Rainfall (mm) | 0.1 | 0.1 | 0 | 0 | 0 |
| Maximum temperature (°C) | 32 | 32 | 32 | 32 | 32 |
| Minimum temperature (°C) | 25 | 25 | 25 | 25 | 25 |
| Maximum Relative Humidity (%) | 74 | 74 | 74 | 74 | 74 |
| Minimum Relative Humidity (%) | 55 | 55 | 55 | 55 | 55 |
| Wind speed (kmph) | 4 | 3 | 2 | 2 | 3 |
| Wind direction (deg) | 270 | 270 | 250 | 270 | 250 |
| Total cloud cover (octa) | 3 | 3 | 2 | 2 | 1 |

Weather summary / Alert: Since dry weather condition is prevailed, irrigation of crops can be continued, when required. Below normal rainfall is expected in Kerala.

B. Agrometeorological Advisories for crop protection & management

C. General and Agrometeorological Advisories for Ernakulum District (RRS Vyttila)

Since there is a chance of dry weather condition due to strong wind and high evaporation, crops should be irrigated, when required. Since the wind speed may increase, give strong propping to banana and staking to vegetables. Proper mulching can be followed to conserve soil moisture.

| Crop | Pest/ Disease | Crop/ Animal Advisory | | |
|--------------------------------------|--|---|--|--|
| Rice (Mundakan) Maturity stage | Bacterial Leaf Blight B | | | |
| Chilly | White fly Since dry weather condition is prevailed, there is a chance of w fly attack in chilly, apply 2% neem oil+ garlic emulsion under leaf surface at 10 days intervals. If the attack is severe, apply 4 g Thiamethoxam per ten litre of water. | | | |
| Arecanut | Spindle bug | There is a chance of spindle bug attack in arecanut plant. The bug sucks the sap from unopened spindle leaf and cause damage. Leaves become dried and shed. As a prophylactic measure, spray 1.5 ml Dimethoate (30 EC) in one litre of water. | | |

Animal Husbandry:- Ensure sufficient supply of clean drinking water to the animals. Allow the animal to graze early in the morning or in the evening to avoid direct sunlight. It is better to graze in the evening itself. (CAADECCS, KVASU, Mannuthy).