

## Kerala Agricultural University, Vellanikkara, Thrissur Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in&kauagmet@yahoo.co.in



(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



## **ERNAKULAM DISTRICT**

AAS 04/2021, (7) Tuesday 19.01.2021

CURRENT SYNOPTIC SITUATION: Northeast monsoon rains have ceased over Tamilnadu, Puducherry & Karaikkal, Kerala & Mahe and adjoining areas of Andhra Pradesh and Karnataka from today, the 19<sup>th</sup> January 2021. The cyclonic circulation over Southwest Bay of Bengal off SriLanka coast lay over Southwest Bay of Bengal and adjoining Equatorial Indian ocean at 5.8 km above mean sea level.

The cyclonic circulation over Equatorial Indian Ocean and adjoining Southwest Bay of Bengal extending upto 2.1 km above me an sea level.

## WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	20.01.2021	21.01.2021	22.01.2021	23.01.2021	24.01.2021
Rainfall (mm)	0	0	0	0	0
Maximum temperature(°C)	32	32	32	32	33
Minimum temperature (°C)	23	23	23	23	24
Maximum Relative Humidity (%)	80	80	80	80	80
Minimum Relative Humidity (%)	40	40	40	40	40
Wind speed (kmph)	4	6	8	6	6
Wind direction (deg)	270	230	320	230	230
Total cloud cover (octa)	2	3	1	3	2

As per the forecast of IMD & KAU, due to no rainfall and prevailing dry weather conditions, irrigation of crops to be continued in the district. Above normal rainfall is expected in Kerala.

## AGRO-METEOROLOGICAL ADVISORIES:

AGRO-METEO	RULUGICAL ADV	ISOKIES.		
Crop	Pest/ Disease	Advisories		
General Advisories: Since no rainfall is forecasted, Drying of agricultural products and spraying of pesticides				
can be done.				
Rice (Mundakan) Dough stage	Bacterial Leaf blight	As a prophylactic measure, spray 20 gram Pseudomonas in one litre of water. To control the spread of Bacterial Leaf blight in paddy, spray 6 g of Streptocycline in 30 litre of water.		
	Sheath Blight	To control sheath blight disease in rice, spray 4 gram of (Trifloxystrobin + Tebuconazole) mixture in ten litres of water.		
	Glume discolouration	During the 50% panicle emergence stage, there is a chance of glume discolouration in rice. To control this, spray 2 gram Kocide per one litre of water.		
Vegetables (Bitter gourd)	Green leaf hopper	There is a chance of green leaf hopper attack in bittergourd. Spray 2 g thiamethoxam per ten litre of water.		
Animal Husbandry	Vaccinate animals against foot and mouth disease and hemorrhagic septicemia. If needed give medicine agaist worm infestation after dung examination.			