



Agromet Advisory Service bulletin 11/17, Tuesday dt 14-03-17 (Issued jointly by IAAS Vellanikkara)





ERNAKULAM

CURRENT SYNOPTIC SITUATION: An upper air cyclonic circulation lies over North Interior Karnataka and neighbourhood upto 1.5 km asl . Another trough extends from this system to extreme south peninsula and extends upto 1.5 km asl.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 14.03.2017 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE: KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 15/03	DAY-2 16/03	DAY-3 17/03	DAY-4 18/03	DAY-5 19/03
ISTRICT : ERNAKULAM					
ainfall (mm)	0	5	2	0	0
ax Temperature (deg C)	33	33	33	33	33
in Temperature (deg C)	25	25	25	25	25
otal cloud cover (octa)	4	4	4	2	2
ax Relative Humidity (%)	85	85	85	80	80
in Relative Humidity (%)	65	65	65	60	60
ind speed (kmph)	800	008	800	800	008
ind direction (deg)	250	250	270	270	270

Agro Meteorological Advisories:

Crop	Stage of the crop / Pest/Disease	Advisories		
Rice		Leaf folder attack can be controlled using trichogramma		
	Tillering stage/Leaf	cards (2CC per one acre). Cut the cards in to small pieces		
	folder	and fix in different regions of the field. If the attack is		
		severe, spray Quinalphos @ 2ml per one liter of water.		
Coconut		Irrigation and mulching is required. If drip irrigation is		
	Management	adopting 60-70 L/day is recommended. Provide adequate		
		shade to the transplanted seedlings.		
Banana		Drench the leaf axils and bore holes on pseudo stem with		
	Pseudostem weevil	Chlorpyrifos 2 ml/L or Quinalphos 2 ml/L if infestation of		
		pseudostem weevil is noticed.		
Ginger/Turmeric	-	Land preparation for planting ginger and turmeric.		
Vegetables (Okra)		Leaves are the most effected plant parts, infected plants		
	Powdery mildew	display white powdery spots on the leaves. Spray		
		karathane 0.5 ml/L in water.		