

Kerala Agricultural University, Vellanikkara, Thrissur **Department of Agricultural Meteorology, College of Horticulture** Email: cohagmet@kau.in&kauagmet@yahoo.co.in **Agromet Advisory Service Bulletin**

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)

ERNAKULAM DISTRICT

30.06.2020

AAS 27/2020,(53) Tuesday CURRENT SYNOPTIC SITUATION: An off-shore trough runs from North Maharashtra to Karnataka coast. A cyclonic circulation lies over Westcentral Arabian sea & neighbourhood between 3.1 km & 3.6 km above mean sea level. A cyclonic circulation lies over North Interior Karnataka & neighbourhood between 4.5 km & 5.8 km above mean sea level.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	01.07.2020	02.07.2020	03.07.2020	04.07.2020	05.07.2020
Rainfall (mm)	2	10	15	25	18
Maximum temperature(°C)	30	30	30	30	30
Minimum temperature (°C)	24	24	24	24	24
Maximum Relative Humidity (%)	96	96	96	96	96
Minimum Relative Humidity (%)	78	78	78	78	78
Wind speed (kmph)	12	12	24	16	20
Wind direction (deg)	270	290	290	320	320
Total cloud cover (octa)	8	8	8	8	8

As per the forecast of IMD & KAU, there is a chance of moderate rainfall in the district.

AGRO-METEOROLOGICAL ADVISORIES: Pest/ Disease

Crop

Advisories

General Advisories: Proper drainage facilities should be ensured to protect the agricultural crops from pest & diseases during rainy season.

Paddy	Seedling stage	During rainy season, the micronutrient deficiency of paddy can be managed by spraying 10 gram of paddy sampoorna in one litter of water.		
Coconut	Management	During the begining of southwest monsoon season, one- kilogram lime can be applid per one matured palm by opening the coconut basin.		
Banana	Pseudo stem weevil	There is a chance of pseudo stem weevil attack in banana. <i>A</i> a precaution, apply 50 g neem cake powder in leaf axils.		
Vegetables	-	During rainy season, the micronutrient deficiency of vegetable can be managed by spraying 5 gram of vegetable sampoorna i one litter of water.		
Animal Husbandry	Keep the cattle shed and premises clean to avoid mosquito breeding. Avoid waterlogging near the cattle shed. The cowdung pit can be covered to protect from direct falling of rain.			