





Agromet Advisory Bulletin for Ernakulam District (Valid from 03.09.2025 to 07.09.2025)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 35/2025, (70) Tuesday

Dated 02.09.2025

A Medium range weather forecast of IMD for next five days

| Weather parameters | 03.09.2025 | 04.09.2025 | 05.09.2025 | 06.09.2025 | 07.09.2025 |
|-------------------------------|------------|------------|------------|------------|------------|
| Rainfall (mm) | 4 | 3 | 2 | 1 | 1 |
| Maximum temperature (°C) | 30 | 30 | 30 | 30 | 30 |
| Minimum temperature (°C) | 25 | 25 | 25 | 25 | 25 |
| Maximum Relative Humidity (%) | 90 | 90 | 90 | 90 | 90 |
| Minimum Relative Humidity (%) | 80 | 80 | 80 | 80 | 80 |
| Wind speed (kmph) | 8 | 4 | 3 | 4 | 8 |
| Wind direction (deg) | 340 | 290 | 290 | 320 | 290 |
| Total cloud cover (octa) | 8 | 8 | 8 | 8 | 7 |

Weather summary / Alert: -. As per the forecast of IMD, there is a chance of light rainfall in the district. Normal to Above Normal Rainfall is expected in Kerala.

B. Agrometeorological Advisories for crop protection & management

C. General and Agrometeorological Advisories for Ernakulum District (RRS Vyttila)

General Advisory: -Special crop management is required for agricultural crops during monsoon season. Proper drainage facilities have to be ensured. Provide supports for the bananas. Make the vegetable pandals stronger. Pesticides should not be applied during rainy period.

| Crop | Pest/ Disease, Crop Management | Agromet Advisory | | |
|--------------------------------|-----------------------------------|--|--|--|
| Paddy (Pokkali) Milky stage | Stem borer | Release of egg parasitoids <i>Trichogramma japonicum</i> (5cc / Ha) at weekly intervals from 15-30 days after transplantation immediately after noticing stem borer activity. | | |
| Paddy (Virippu) Milky stage | Rice bug | Fish amino acid @ 20 ml /L water or Azadirachtin (3000 ppm) 10 ml / L water can be sprayed. Farmers those who cultivates for second crop can initiate preparations for nursey. | | |
| Coconut | Bud rot | Ensure drainage. Prophylatic management practices against bud rot to be taken | | |
| l (inger | | Spray <i>Beauveria bassiana</i> and <i>Pseudomonas</i> @ 20g in a litre of water in alternate weeks | | |

Fisheries: - Take precautions to avoid ectoparasitic infestation. If fever, shivering, brown coloured urine, off feed etc. are noticed, consult veterinary doctor immediately. Ensure sufficient supply of clean drinking water to the animals.