


Agromet Advisory Service bulletin 41/17, Tuesday dt .10.10.2017





Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara E mail: kauagmet@yahoo.co.in, cohagmet@kau.in

ERNAKULAM

CURRENT SYNOPTIC SITUATION: A Deep Depression lies over Gangetic West Bengal near Lat 23.5°N and Lon 86.7°E. An upper air cyclonic circulation lies over East central Arabian sea and neighbourhood and extends upto 3.6km above sea level.

Heavy Rainfall Warning: Heavy rainfall is most likely to occur at one or two places in Kerala from 11th till the morning of 13th October 2017.

> NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 10.10.2017 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

DISTRICT: ERNAKULAM ENSEMBLE FCST

PARAMETERS	11.10.2017	12.10.2017	13.10.2017	14.10.2017	15.10.2017
Rainfall (mm)	3	22	23	22	18
Max Temperature (deg	32	32	32	31	31
Min Temperature (deg C)	25	25	25	24	24
Total cloud cover (octa)	5	8	8	8	8
Max Relative Humidity (%)	90	90	90	90	90
Min Relative Humidity (%)	80	80	80	80	80
Wind speed (kmph)	14	14	14	14	14
Wind direction (deg)	180	180	250	250	250

Agro Meteorological Advisories:

Crop	Crop stage / Pest/Disease	Advisories	
Paddy	Harvesting	First paddy crop is in Harvesting stage. Land preparation for second	
(Jyothi, Kanchana)	stage	paddy crop can be started.	
		As a prophylactic measure, hung three perforated packets of	
Coconut	Bud rot	Mancozeb sachet (5gm) around spindle leaves.	
		Destroy the affected leafs along with the spodoptra. For severe attack	
Banana	Spodoptra	spray Flubendiamide (Fame) 2 ml/10 L of water.	
		This is the best time to sow cabbage and cauliflower seeds in the nursery beds. Addition of pseudomonas culture to the soil at the time	
Vegetables	-	of planting of cabbage and cauliflower will help to reduce fungal	
	-	diseases.	
	Deworming of animals should be done only after dung sample examination. In calves		
Animal Husbandry	first dose of medicine should be given within 15 days of age. (CAADECS, KVASU, Mannuthy)		