

Agromet Advisory Bulletin for the District, Kasaragod (Valid from 28.10.2023to 01.11.2023)



(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode& India Meteorological Department)

Bulletin Number:Pilicode/Ksd-86/2023 Date:27/10/2023
--

A. Weather Summary of preceding four days

Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
0.0	33.0 – 33.5	24.5 – 25.2	73–92	01 - 03

B. Weather forecast for next five days

Parameters	28-10-2023	29-10-2023	30-10-2023	31-10-2023	01-11-2023
Rainfall, mm	0.5	6	3	5	1
Max. Temp, °C	34	34	34	34	34
Min. Temp,°C	24	24	24	24	24
Max. Relative Humidity, %	84	86	88	90	86
Min. Relative Humidity, %	70	72	74	75	78
Wind speed,km/h	2	2	2	2	2
Wind direction, degrees	230	290	290	270	250
Total cloud cover, octa	7	8	8	8	8

C. Agrometeorological Advisories

Crop	Stages	Problems	Agro-meteorological advisories	
	Light to Moderate rainfalls **			
General	The sky will be cloudy. High relative humidity will be experienced.			
conditions	Low night temperature and high day temperature will be experienced. Hence there be distinct difference between day time temperature and night temperature.			
	There will be light rainfalls (from 2.5mm to 15.5 mm within a time span of 24 hours) in most places from October 27 th to October 31 th .			
General Recommen	The weather is favorable for the spread of fungal diseases like bud rot of coconut, mahali in arecanut, quick wilt in black pepper, rhizome rot diseases in ginger and turmeric.			
dations	Maintain hygiene conditions in crop fields. Infected and fallen nuts, leaves and tree parts should be removed from the fields and burnt.			
	Provide propping to tender stem crops like banana, tomato etc. Ensure sufficient drainage in crop fields. Beware of lightning. Give popping to slender stemmed and easily lodging crops like banana, vegetables etc. Take care while drying the harvested produces like rubber, cashew nut, copra etc., directly under the sun. Beware of lightning.			
	To cop up the summer season of 2024 year, the preparatory measures have to be undertaken from now onwards. As rainfall is predicted, ensure maximum water harvest in the fields. Clean the			

	which are partially fill	l with dried leaves, especially the basins of crops. The opened tree basins led after fertilizer application, can act as very good water harvesting the runoff water to such tree basins by drawing furrows.		
Vegetables	Transplanting/ sowing	While preparing the land, incorporate lime @ 4kg/cent to the soil. Use Trichoderma enriched farm yard manure/compost. This will check the spread of wilt diseases. Before transplanting, dip the roots of the seedlings in slurry of pseudomonas (20g pseudomonas/litre of water) for 30 minutes. This will help the seedlings to grow vigorously. Also the seedlings can be sprayed with diluted pseudomonas culture solution (@20ml dissolved in one litre of water).		
Coconut	All stages	Red palm weevil	The palms can be saved only if the attack is diagnosed at an early stage. Hence more frequently inspect the palms. The attack can be diagnosed by the presence of yellowing of upper and middle whirls of leaves and holes at the jointing region of leaf petioles to the trunk. Insect excrement, pupal cases etc. may also be seen in the crown or basins of the palms.	
			Control: Prepare Imidachlorprid solution (@ 1ml/litre of water) and fill in the holes. Adopt field sanitation.	
Black pepper	All stages	Foot rot	As prophylactic measure, apply 150 gram of Trichoderma enriched neem cake - cow dung mixture in the basins of the vines and incorporate thoroughly with the soil. If disease already appeared, drench soil in the plant basins with Redomil 0.2% (2g/litre of water). Spray the same on the leaves also.	
Banana	Various stages of growth	Sigatoka leaf spot	There should not be any water stagnation in the basins of the crop. As a prophylactic measure spray Pseudomonas (@ 20g/litre) at fortnight intervals.	
			If disease appears:-Cut and burn all affected leaves. Spray Carbendazim and Mancozeb (@ 2ml per litre) alternatively at fortnight intervals	
Cashew	Pre-bearing stages	Tea mosquito bug	Prophylactic measures:	
			1) The fungal biological control agent, Beavaria bassiana can be sprayed @ 20g/L Or	
			2) Spray 1% Bordeaux mixture mixed with quinalphos (2ml/litre of Bordeaux mixture)	

Ginger	All stages	Fusarium wilt	Drain the field. Drench the field with copper oxy chloride solution (@2g/litre of water)
Turmeric	Different stages	Leaf blight	Remove the severely affected leaves from the plants and destroy in fire. Spray copper oxy chloride on the plants (@2g/litre of water)
Vegetables	Planting	Downy mildew on cucurbits	As a prophylactic measure apply 'Mancozeb' (@ 2g/l of water). If disease appeared, spray Akomin® (@3ml/L) on both surfaces of the leaves, thrice at 15 days interval. Remove and burn out the infested leaves
Cow	Any stage	FMD (Foot and Mouth disease)	Foot and Mouth disease is a highly transmissible disease caused by infection with an Aphthovirus. The infection results in vesicular lesions in and around the mouth and on the feet, resulting in the reluctance of an animal to eat or move.
			Prevention: Ensure hygienic conditions for the individual animals as well as their shelter and its surroundings. Boost the immunity of cattle by supplementing feeds with vitamins and minerals containing health tonics.
** Warning ca	olour codes of rainfall (for di	saster management)	If disease appears, give medication under the supervision of a Veterinary doctor immediately.

Warning (Take actions)

Alert (Be prepared)

No warning (No actions)

Watch (Be updated)