

Kerala Agricultural University, Vellanikkara, Thrissur Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in & kauagmet@yahoo.co.in







OLLUKKARA BLOCK, THRISSUR DISTRICT

AAS 46/2018(90), Tuesday

13.11.2018

The Agricultural advisory of this bulletin is not applicable in flood-affected areas.

SIGNIFICANT WEATHER FOR THE PRECEDING WEEK AT VELLANIKKARA:

The total rainfall received 0.0 mm. the maximum temperature ranges from 33.8°C to 34.0°C, minimum temperature ranges from 21.9°C to 23.4°C. Morning relative humidity ranges from 77% to 90% and afternoon relative humidity ranges from 40% to 57%. Wind speed ranges from 2.1 to 3.8 km/h. Bright sunshine hours range from 4.3 to 9.9 hours. Evaporation ranges from 3.1 to 4.0 mm.

CURRENT SYNOPTIC SITUATION: The cyclonic storm 'Gaja' lay centered at 0830 hrs. IST over west central and adjoining East central & South west Bay of Bengal near Lat 13.3° N / long 87.1° E about 740 km east Northeast of Chennai

HEAVY RAINFALL WARNING: DAY 3: Heavy to very heavy rainfall is most likely to occur in Kollam District and heavy rainfall is most likely to occur at isolated places in Trivandrum & Pathanamthitta district.

DAY 4: heavy to very heavy rainfall is most likely to occur at isolated places in Trivandrum & Kollam district and heavy rainfall is most likely to occur at isolated places in Pathanamthiita & Alappuzha district.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	14.11.2018	15.11.2018	16.11.2018	17.11.2018	18.11.2018
Rainfall (mm)	4	2	0	25	10
Maximum temperature (°C)	32.8	33.2	34	34	33.8
Minimum temperature (°C)	22.8	23	23.8	23.8	23.8
Total cloud cover (octa)	5	5	6	7	7
Maximum Relative Humidity (%)	90	85	75	90	90
Minimum Relative Humidity (%)	60	60	55	60	60
Wind speed (kmph)	6	8	6	10	8
Wind direction (deg)	320	360	50	50	90
A TMTD C 4 12-1-4 2 C-11 2-	4 1 6	. 1 . 41	11 1		

As per IMD forecast, light rainfall is expected for coming days in the block.

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Crop stage	Pest/ Disease	Advisories
Paddy	Seedling stage	Rice armyworm	 There is a chance of rice armyworm in the paddy field. To control, Complete flooding of field for 24 hours wherever possible. Spraying of chlorantraniliprole @ 3 ml per 10 litre of water in the field, bunds and weeds. In the fields where transplanting is not yet undertaken, application of chlorantraniliprole granules @ 4 kg per one acre along with basal dose of fertilizers. Keep the field bunds and surroundings weed free.
Banana	-	Spodoptera	Destroy the affected leafs along with the <i>Spodoptera</i> . If <i>Spodoptera</i> attack is severe, spray quinalphos 2ml per one litre of water or chlorantraniliprole 3 ml per 10 litre of water.
Pepper	-	Yellowing	Yellowing was observed in pepper due to flood situation followed by physiological drought. Soil drenching with 2g kocide per one litre of water,

followed by a foliar spray and soil drenching with

	20 g of pseudomonas per litre of water, after two weeks i/s recommended.				
Animal	Mix 30g sodium bicarbonate and 1-teaspoon yeast in the feed daily to avoid digestive				
Husbandry	disorders associated with excessive acidity. (CAADECCS, KVASU, Mannuthy)				
	,				