



**Kerala Agricultural University, Vellanikkara, Thrissur**  
**Department of Agricultural Meteorology, College of Horticulture**

Email: cohagmet@kau.in&kauagmet@yahoo.co.in

**Agromet Advisory Service Bulletin**

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



**OLLUKKARA BLOCK, THRISSUR DISTRICT**

**AAS 44/2020,(88) Friday**

**30.10.2020**

**CURRENT SYNOPTIC SITUATION:** A cyclonic circulation lies over Tamil Nadu coast & adjoining Southwest Bay of Bengal extending upto 4.5 km above mean sea level. A cyclonic circulation lies over Eastcentral Bay of Bengal & adjoining North Andaman Sea between 1.5 & 4.5 km above mean sea level. A cyclonic circulation lies over South & adjoining Central Arabian Sea extending upto 1.5 km above mean sea level persists..

**WEATHER FORECAST FOR THE NEXT FIVE DAYS:**

Weather parameters	31.10.2020	01.11.2020	02.11.2020	03.11.2020	04.11.2020
Rainfall (mm)	0.9	0	0	0	18.3
Maximum temperature(°C )	35.7	34.4	34.7	33.9	31.1
Minimum temperature (°C )	23.5	23.9	23.9	24.6	24.5
Maximum Relative Humidity (%)	93	95	96	94	96
Minimum Relative Humidity (%)	46	49	51	52	80
Wind speed (kmph)	4	4	4	5	4
Wind direction (deg)	108	108	68	68	72
Total cloud cover (octa)	5	7	4	7	8

As per the forecast of IMD & KAU, there is a chance of low rainfall in the district.

**AGRO-METEOROLOGICAL ADVISORIES:**

Crop	Pest/ Disease	Advisories
<b>General Advisories:</b> Since low rainfall is expected, spraying of pesticides and drying of agricultural products can be done.		
Rice (Mundakan) Seedling stage	Leaf folder	Since cloudy weather is prevailed, there is a chance of leaf folder attack in <i>Mundakan</i> Paddy. It can be controlled by using <i>trichogramma</i> cards (2CC per one acre). Cut the cards in to small pieces and fix in different regions of the field. If the attack is severe, spray quinalphos @ 2ml per one litre of water.
	Army worm	There is a chance of rice armyworm in the paddy field. To control, <ul style="list-style-type: none"> <li>• Complete flooding of field for 24 hours wherever possible.</li> <li>• Spraying of chlorantraniliprole @ 3 ml per ten litre of water in the field, bunds and weeds.</li> <li>• In the fields where transplanting is not yet undertaken, application of chlorantraniliprole granules @ 4 kg per one acre along with basal dose of fertilizers.</li> <li>• Keep the field bunds and surroundings weed free.</li> </ul>
Banana	Planting time	Lime can be applied at 250 gram per plant at the time of planting.
Brinjal	White fly	There is a chance of white fly attack in Brinjal. Apply 2% neem oil garlic emulsion or spray 20g <i>Lecanicillium lecanii</i> per one litre of water in ten days intervals.
Animal Husbandry		Mix 30g sodium bicarbonate and 1-teaspoon yeast in the feed daily to avoid digestive disorders associated with excessive acidity.