

Agromet Advisory Service bulletin 23/17 Tuesday dt 06.06.2017 (Issued jointly by IAAS Vellanikkara) Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara E mail: kauagmet@yahoo.co.in, cohagmet@kau.

PALAKKAD

CURRENT SYNOPTIC SITUATION: A well marked low pressure area lies over west central Arabian sea with associated upper air cyclonic circulation extending upto mid tropospheric levels. A trough at mean sea level runs from south Maharashtra coast to Kerala coast. An upper air cyclonic circulation lies over north Andhra Pradesh coast and neighbourhood between 3.1 & 3.6 km above mean sea level.

HEAVY RAINFALL WARNING: Heavy rainfall will occur at one or two places in Kerala till the morning of 8th June 2017.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 06.06.2017 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE: KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 07/06	DAY-2 08/06	DAY-3 09/06	DAY-4 10/06	DAY-5 11/06
DISTRICT : PALAKKAD					
Rainfall (mm)	25	20	15	20	15
Max Temperature (deg C)	31	31	31	31	31
Min Temperature (deg C)	24	24	24	24	24
Total cloud cover (octa)	8	7	7	7	7
Max Relative Humidity (%)	90	90	90	90	90
Min Relative Humidity (%)	80	80	80	80	80
Wind speed (kmph)	012	012	012	012	012
Wind direction (deg)	250	250	250	250	250

Agro Meteorological Advisories:

Сгор	Stage of the crop / Pest/Disease	Advisories	
Paddy	Land preparation to transplanting	Transplant seedlings at 4-5 leaf stage. Soak seed for 30 minutes	
		in a solution of Pseudomonas culture @10g/litre per kg of	
		seeds. The above seed treatment can also be used to control	
		sheath blight, sheath rot, leaf spot disease etc.	
Banana	Sigatoka disease	Due to the pre monsoon showers, there is chance of Sigatoka	
		disease in banana. As a precaution, Spray Pseudomonas 20g/li.	
		In case of severe attack spray 1 gm Bavistin by mixing it with	
		gum in 1 litre of water on lower surface of leaf.	
Vegetables	Powdery mildew	Threre is a chance of powdery mildew attack in vegetables.	
		Application of mancozeb 3g/L or Trichoderma 2g/L is	
		effective.	