



Agromet Advisory Service bulletin 30/17 Tuesday dt 25.07.2017

(Issued jointly by IAAS Vellanikkara)





PALAKKAD

CURRENT SYNOPTIC SITUATION: A well marked low pressure area lies over Gangetic West Bengal & adjoining Jharkhand with associated upper air cyclonic circulation extending up to 9.5 km above sea level. An off shore trough at mean sea level runs from South Gujarat coast to north Kerala coast.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 25.07.2017 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE: KERALA

PARAMETERS	ENSEMBLE FCST				
	ENSEMBLE FCS1				
	DAY-1	DAY-2	DAY-3	DAY-4	DAY-5
	26/07	27/07	28/07	29/07	30/07
DISTRICT : PALAKKAD					
Rainfall (mm)	2	2	5	6	10
Max Temperature (deg C)	32	32	32	32	32
Min Temperature (deg C)	24	24	24	24	24
Total cloud cover (octa)	5	5	5	6	6
Max Relative Humidity (%)	90	90	90	90	90
Min Relative Humidity (%)	80	80	80	80	80
Wind speed (kmph)	012	012	012	012	012
Wind direction (deg)	250	250	250	250	250

Agro Meteorological Advisories:

Crop	Stage of the crop / Pest/Disease	Advisories		
D 11	D : 1	Chances of stem borer in paddy. Use 2 CC trichocards as		
Paddy (Jyothi, Kanchana)	Panicle emergence/ stem borer	10 pieces in different corners of field under leaf. In severe		
	Stelli bolei	condition apply Flubendamide in 2 ml/10 L		
Banana		Due to the pre monsoon showers, there is chance of		
		Sigatoka disease in banana. As a precaution, Spray		
	Sigatoka disease	Pseudomonas 20g/L In case of severe attack spray 1 g		
		Bavistin by mixing it with gum in 1 litre of water on lower		
		surface of leaf.		
Ginger, Turmeric	Rhizome rot	During rainy season there is a chance of rhizome rot, to control this dig out the affected plants and apply Kocide 2gm/1 litre water.		
		Take vaccination against Haemorrhagic septicaemia in		
		cattles.		
Animal Husbandry				
	-	Thank you for Information: Centre for Animal		
		Adaptation Environment and Climate Change Studies,		
		KVASU, Mannuthy.		