11/14/25, 3:24 PM District Advisory



Gramin Krishi Mausam Sewa Agromet Advisory Bulletin for THRISSUR District Kerala Agricultural University



Vellanikkara, Thrissur, Kerala

Agromet Advisory Bulletin

Date: 2025-11-14

Weather Forecast of District THRISSUR (Kerala) Issued On: 2025-11-14 (Valid Till 08:30 IST of the next 5 days)

Parameter	2025-11-15	2025-11- 16	2025-11- 17	2025-11-18	2025-11-19
Rainfall(mm)	0.5	0.1	3.0	0.1	0.3
Tmax(°C)	34.0	34.0	34.0	34.0	34.0
Tmin(°C)	23.0	23.0	23.0	23.0	23.0
RH-I(%)	80	80	80	80	80
RH-II(%)	70	70	70	70	70
Wind Speed(kmph)	2	2	2	2	4
Wind Direction(Degree)	270	230	270	20	50
Cloud Cover(Octa)	8	5	8	8	7
Warning	Heavy Rain; Thunderstorms & Lightning, Squall etc	No Warning	No Warning	Thunderstorms & Lightning, Squall etc	Thunderstorms & Lightning, Squall etc

Forecast Summary:

Weather summary / Alert:-As per the forecast of IMD, there is a chance of light rainfall in isolated places of the district. Thunderstorm accompanied by lightning is very likely to occur at one or two places in Kerala. Below normal rainfall is expected in Kerala.

Weather Warnings (Valid Till 08:30 IST of the next day)

Heavy Rain; Thunderstorms & Lightning, Squall etc

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Thunderstorm accompanied by lightning is very likely to occur at one or two places in Kerala.

General Advisory:

General Advisory: -Prophylactic measures can be taken to control crop pest & diseases. Proper mulching of crops can be followed to conserve soil moisture.

SMS Advisory:

As per the forecast of IMD, there is a chance of light rainfall in isolated places of the district.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory		
	Rice(Mundakan)-Tillering stage- Thrips -To control thrips in rice, spray 2 gram of Thiomethoxam in 10 litre of water.		

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory	
OKRA/ LADYFINGER (Flowering)	Bhindi - Jassid's -Due to increase in atmospheric temperature, the attack of Jassid's in Bhindi was increased. They can be controlled by applying 2% neem oil emulsion once in a week or by spraying 2 g thiamethoxam per ten litre of water.	
	Cowpea- Web blight -There is a chance of web blight disease in cowpea. As prophylactic measure soil drenching of cowdung slurry supernatant 2% + Pseudomonas fluorescens 2% at 20, 40 and 60 days after sowing.	

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory		
COW	Animal Husbandry: - Precautions should be taken to avoid ectoparasitic infestations. Deworming of animals should be done after microscopical examination of dung sample in a veterinary hospital. (CAADECCS, KVASU, Mannuthy).		

Likely impacts of weather warnings (General)

Harvest the crop as early as possible. Provide proper drainage in the cropped field as heavy rainfall is expected. Propping of the banana plants recommended. Additional support of the pandals recommended. Keep away from open fields to avoid lightning stroke to livestock and farmers in the afternoon.Livestock / Poultry / Fishery: Keep the animals inside the shed during heavy rainfall / hailstorm and provide them balanced feed.

Impact based advisories (General)

Heavy rains may cause flooding in the lowlands. Heavy rainfall results in the flooding of the cropped field which damages the crop in the maturity stage. High winds cause lodging of the banana plants. Those vegetables in pandals will be damaged due to high winds.

Farmers are advised to download Unified "Mausam" and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

11/14/25, 3:24 PM District Advisory

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link: https://play.google.com/store/apps/details