







Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 01/2024, (2) Friday

Dated 05.01.2024

	Observed 0 th Decer				2024)		Weather Forecast Valid for 06 ^h January to 10 th January 2023						
30/12	31/12	01/01	02/01	03/01	04/01	05/01	Date		06/01	07/01	08/01	09/01	10/01
0.0	0.0	0.0	0.0	0.0	0.0	94.0	Rainfall (mm)		11	11	21	2	0.2
32.8	31.7	32.0	31.9	32.1	31.2	32.4	T _{max} (°C)		33	33	33	33	33
25.4	24.6	24.4	24.3	24.4	24.6	23.3	T min(⁰ C)		23	24	24	27	27
Cloudy	Partially cloudy	Partially cloudy	Cloudy	Partially cloudy	Cloudy	Cloudy	AM	Cloud	Overcast	Overcast	Overcast	Overcast	Overcast
Partially cloudy	Partially cloudy	Cloudy	Cloudy	Cloudy	Overcast		PM	cover					
75	74	80	76	79	79	90	RH-I (%)		74	74	74	74	74
54	51	57	52	57	56		RH-II (%)		55	55	55	55	55
11.5	11.8	8.8	6.8	8.2	5.4	6.2	Wind Speed (km/hr.)		3	2	2	2	2
NNE	NE	NNE	NNE	NNE	NNE	ENE	AM	Wind	WNW	SW	W	wsw	WNW
SE	NE	NE	NNE	NE	NE		PM	direction					
Rainfall (mm) in last week Rainfall					fall (mm	(mm) from 01/01/2024 to till dated				Rainfall (mm) from 01/10/2024 to today			
94.0				94.0					94.0				

Weather summary / Alert: As per the forecast of IMD, there is a chance of moderate rainfall in the district. Above normal rainfall is expected in Kerala.

C. General and Agrometeorological Advisories for Thrissur District

Since there is a chance of dry weather condition due to strong wind and high evaporation, crops should be irrigated, when required. Since the wind speed may increase, give strong propping to banana and staking to vegetables. Proper mulching can be followed to conserve soil moisture.

bCrop	Pest/ Disease	Crop/ Animal Advisory					
Rice (Mundakan) Panicle stage	-	In order to reduce the micronutrient deficiency in paddy, spray 10 grams of sampoorna mixture of paddy in one liter of water. Sampoorna mixture of paddy is available at the Kerala Agricultural University.					
Banana	-	30 grams of urea and 60 grams of potash can be applied to each banana plant of four months old.					
Tomato	White fly	Since dry weather condition is prevailed, there is a chance of white fly attack in tomato, apply 2% neem oil+ garlic emulsion under the leaf surface at 10 days intervals. If the attack is severe, apply 4 gram Thiamethoxam per ten litre of water.					

Animal Husbandry:-Vaccinate animals against foot and mouth disease and hemorrhagic septicemia. If needed give medicine agaist worm infestation after dung examination. (CAADECCS, KVASU.