



Kerala Agricultural University, Vellanikkara, Thrissur
Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in&kauagmet@yahoo.co.in

Agromet Advisory Service Bulletin

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



THRISSUR DISTRICT

AAS 06/2021, (11) Tuesday

02.02.2021

SIGNIFICANT WEATHER FOR THE PRECEDING WEEK AT VELLANIKKARA:

The total rainfall received 0.0 mm. The maximum temperature ranges from 32.7°C to 34.2°C, minimum temperature ranges from 18.1°C to 21.7°C. Morning relative humidity ranges from 58% to 85% and afternoon relative humidity ranges from 31% to 40%. Wind speed ranges from 3.4 to 6.5 km/h. Bright sunshine hours range from 8.0 to 10.0 hours. Evaporation ranges from 4.4 to 5.8 mm.

CURRENTSYNOPTICSITUATION:

A trough in low level easterlies runs from north Kerala coast to south Gujarat coast upto 0.9 km above mean sea level.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	03.02.2021	04.02.2021	05.02.2021	06.02.2021	07.02.2021
Rainfall (mm)	0	0	0	0	0
Maximum temperature(°C)	34	34	34	34	34
Minimum temperature (°C)	22	22	22	22	22
Maximum Relative Humidity (%)	67	67	67	67	67
Minimum Relative Humidity (%)	36	36	36	36	36
Wind speed (kmph)	20	18	20	4	2
Wind direction (deg)	50	50	50	50	20
Total cloud cover (octa)	2	3	4	3	3

As per the forecast of IMD & KAU, due to no rainfall and prevailing dry weather conditions, irrigation of crops to be continued in the district. Below normal rainfall is expected in Kerala

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Pest/ Disease	Advisories
General Advisories: Since no rainfall is forecasted, drying of agricultural products and spraying of pesticides can be done.		
Rice (Mundakan) Dough stage	Bacterial Leaf Blight	As a prophylactic measure, spray 20 gram Pseudomonas in one litre of water. To control the spread of Bacterial Leaf blight in paddy, spray 6 g of Streptocycline in 30 litre of water.
	Blast	To control Blast disease in rice, spray 4 gram of Nativo (Trifloxystrobin + Tebuconazole) mixture in ten litres of water.
Banana	Boron deficiency	Boron deficiency is seen more during dry weather. Spray one gram borax per one litre of water.
Cucurbitaceous vegetables	Fruit fly	Remove and destroy affected and decayed fruits. Use special pheromone traps like cuelure (6 no./acre) or Thulasi/fruit traps for vegetable fruit fly. If the attack is not reduced, spray 2 ml Cyantraniliprole in one litre of water
Animal Husbandry		Ensure sufficient supply of clean drinking water to the animals. Take measures for preventing mosquito breeding in the premises.