




## Advisory Service bulletin 24/17, Friday dt.16-06-2017

( Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara

E mail: kauagmet@yahoo.co.in, cohagmet@kau.in



### THRISSUR

**CURRENT SYNOPTIC SITUATION:** An off shore trough runs from south Maharashtra coast to Kerala coast.

**HEAVY RAINFALL WARNING:** Heavy rainfall will occur at one or two places in Kerala from the morning of 18<sup>th</sup> to 20<sup>th</sup> June 2017.

#### NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST

ISSUED ON: 16.06.2017

VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE: KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 17/06	DAY-2 18/06	DAY-3 19/06	DAY-4 20/06	DAY-5 21/06
DISTRICT : THRISSUR					
Rainfall (mm)	11	3	15	40	19
Max Temperature ( deg C)	32	32	32	30	30
Min Temperature ( deg C)	25	24	24	24	24
Total cloud cover (octa)	7	6	7	8	8
Max Relative Humidity (%)	90	90	90	90	90
Min Relative Humidity (%)	80	80	80	80	80
Wind speed (kmph)	010	010	010	010	010
Wind direction (deg)	250	250	250	250	250

#### Agro Meteorological Advisories:

Crop	Stage of the crop / Pest/Disease	Advisories
Paddy	Land preparation to transplanting	Transplant seedlings at 4-5 leaf stage. Soak seed for 30 minutes in a solution of Pseudomonas culture @10g/litre per kg of seeds. The above seed treatment can also be used to control sheath blight, sheath rot, leaf spot disease etc.
Banana	Rhizome rot	Apply pseudomonas fluorescens 20g/L to control the rhizome rot. For one banana plant 4 L solution can be applied.
Vegetables	Powdery mildew	There is a chance of powdery mildew attack in vegetables. Application of mancozeb 3g/L or Trichoderma 20g/L is effective.