



## Agromet Advisory Service bulletin 37/17, Tuesday dt 12.09.2017

(Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara E mail: kauagmet@yahoo.co.in, cohagmet@kau.in







**CURRENT SYNOPTIC SITUATION:** An upper air cyclonic circulation lies over south east Arabian sea off Lakshadweep area and extends upto 2.1 km above sea level.

## NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 12.09.2017 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

DISTRICT: THRISSUR ENSEMBLE FCST

PARAMETERS	13.09.2017	14.09.2017	15.09.2017	16.09.2017	17.09.2017
Rainfall (mm)	16	15	23	13	20
Max Temperature ( deg	33	33	31	30	30
Min Temperature ( deg C)	24	24	24	24	24
Total cloud cover (octa)	7	7	8	7	8
Max Relative Humidity (%)	90	90	90	90	90
Min Relative Humidity (%)	80	80	80	80	80
Wind speed (kmph)	10	10	10	10	10
Wind direction (deg)	180	180	250	270	270

## **Agro Meteorological Advisories:**

Crop	Stage of the crop / Pest/Disease	Advisories	
		As a prophylactic measure, hung three perforated packets	
Coconut	Bud rot	of Mancozeb sachet (5gm) around spindle leaves.	
		Due to continuous rain, there is a chance of Sigatoka	
		disease in banana. As a precaution, Spray Pseudomonas	
Banana	Sigatoka disease	20g/litre. In case of severe attack, spray 1 gm of Bavistin	
		in 1 litre of water on lower surface of leaf (by mixing with	
		gum)	
		If yellowing symptom due to downy mildew is observed in	
Vegetables	Downy mildew	cucurbitaceous vegetables, apply 3 g/Mancozeb per liter of water.	
		Mosquito control measures can be adopted. Cattle should	
		not be allowed graze in water logged areas for avoiding	
		diseases. Avoid water logging in cattle sheds also. If fever	
Animal Husbandry	-	and respiratory distress are noticed consult a veterinary	
		surgeon.(Centre for animal adaptation to environment and	
		climate change studies, KVASU, Mannuthy)	