



Agromet Advisory Bulletin for Thrissur District





Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 38/2023, (76) Friday

Dated 22.09.2023

(Dated	Observed 16 th Sept		U	et week ptember 20	23)		Weather Parameters		Weather Forecast Valid for 23 rd September to 27 th September 2023)				
16/09	17/09	18/09	19/09	20/09	21/09	22/09	Date		23/09	24/09	25/09	26/09	27/09
23.0	19.7	14.6	23.0	0.2	0.0	2.6	Rainfall (mm)		4	2	11	11	5
29.2	28.2	30.7	30.0	31.6	31.5	31.9	T _{max} (⁰ C)		32	33	33	33	33
23.4	23.7	24.5	23.3	24.4	24.3	23.9	T min(⁰ C)		24	25	25	25	25
Cloudy	Overcast	Overcast	Partially cloudy	Cloudy	Partially cloudy	Partially cloudy	AM	Cloud	Overcast	Overcast	Overcast	Overcast	Overcast
Cloudy	Cloudy	Cloudy	Cloudy	Partially cloudy	Cloudy		PM	cover					
99	97	99	96	93	92	95	RH-I (%)		93	94	94	95	94
89	78	81	68	71	65		RH-II (%)		66	68	60	63	63
1.2	0.7	1.1	0.6	15	1.0	1.1	Wind Speed (km/hr.)		2	3	4	4	3
C	С	C	C	C	C	C	AM	Wind direction	WNW	W	NW	WNW	W
SW	SW	WNW	WSW	SW	NW		PM						
Ra	Rainfall (mm) in last week Rainfall (mm) from 0					1/01/2023 t	o till dated	Rainfall (mm) from 01/06/2023 to today					
83.1				1419.6					1286.2				

Weather summary / Alert: As per the forecast of IMD, there is a chance of light rainfall in the district and moderate rainfall in isolated places. Above normal rainfall is expected in Kerala.

C. General and Agrometeorological Advisories for Thrissur District

Prophylactic measures to control crop pest & diseases can be taken to avoid their spread. Proper mulching can be followed to conserve soil moisture.

Crop	Pest/ Disease	Crop/ Animal Advisory
Paddy (virippu season) Panicle stage	Rice bug	There is a chance of appearance of rice bug in paddy. To control, spray 2 ml malathion per one litre of water. Pesticides application should be done either before 9 am or after 3 pm. Otherwise, it will adversely affect the pollination.
Pepper	Quick wilt	As a prophylactic measure to control quick wilt of pepper, add 2 kg Trichoderma to a mixture of 90 kg FYM and 10 kg neem cake and kept for two weeks with sufficient moisture. Add 2.5 kg of this mixture to the base of each matured pepper vine
Ginger	Stem borer	There is a chance of stem borer in ginger. Spary 20 g <i>Beauveria per</i> one litre of water to control stem borer. Or spray 2 ml of quinalphos in one litre of water.
Cucurbitaceous Vegetables	Downy mildew	There is a chance of downy mildew attack in cucurbitaceous vegetables. Spray 2.5gm Mancozeb per one litre of water under the leaves.

Animal Husbandry: - Take precautions to avoid ectoparasitic infestation. If Fever, shivering, and brown coloured urine or any other symptoms of jaundice if noted, immediately consult a veterinary doctor.