



## Agromet Advisory Bulletin for Thrissur District (Valid from 28.10.2023 to 01.11.2023)



Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

## AAS Bulletin No: 43/2023, (86) Friday

Dated 27.10.2023

(Dated			during las to 27 <sup>th</sup> O	t week ctober 202.	3)		Weather	Parameters	Weather Forecast Valid for 28 <sup>th</sup> October to 01 <sup>st</sup> November 2023)				
21/10	22/10	23/10	24/10	25/10	26/10	27/10	Date		28/10	29/10	30/10	31/10	01/11
0.0	0.4	15.2	0.0	0.0	3.4	4.6	Rainfall (mm)		9	10	8	14	0.5
33.4	33.6	32.5	32.0	33.0	32.5	33.7	T <sub>max</sub> ( <sup>0</sup> C)		34	33	33	33	33
24.6	25.2	23.3	23.4	24.5	24.1	24.5	T min( <sup>0</sup> C)		25	24	24	24	24
Cloudy	Cloudy	Partially Cloudy	Cloudy	Overcast	Partially Cloudy	Partially Cloudy	AM	Cloud	Overcast	Overcast	Cloudy	Overcast	Overcast
Cloudy	Overcast	Overcast	Partially Cloudy	Overcast	Overcast		PM	cover					
95	93	96	95	93	89	93	<b>RH-I</b> (%)		91	92	91	94	93
63	70	75	59	69	55		<b>RH-II</b> (%)		72	70	72	75	70
1.3	1.0	1.2	1.6	1.4	1.1	0.6	Wind Speed (km/hr.)		2	2	2	2	2
С	С	С	С	С	С	С	AM	Wind direction	SW	W	W	WSW	W
С	NNE	SE	С	SW	С		PM						
Rainfall (mm) in last week Rainfall (mm) from 0						l/01/2023 t	o till dated	Rainfall (mm) from 01/10/2023 to today					
23.6 177					1774	.2		155.2					

Weather summary / Alert: As per the forecast of IMD, there is a chance of moderate rainfall in isolated places of the district. Above normal rainfall is expected in Kerala.

## C. General and Agrometeorological Advisories for Thrissur District

Prophylactic measures can be taken to control and to avoid the spread of crop pest and diseases. Proper mulching can be followed to conserve soil moisture.

Сгор	Pest/ Disease	Crop/ Animal Advisory				
Rice		As a prophylactic measure to manage sheath blight, sheath rot,				
(Mundakan) Seedling stage	-	leaf spot disease, spray 20g of pseudomonas in one litre of water.				
		In the second stage of fertilizing coconut plants, can apply 1.5				
Coconut	-	kg of urea, 1.45 kg of rock phosphate and 2.25 kg of muriate of potash per coconut plant.				
		In the first stage of fertilizing arecanut plants, can apply 109				
Arecanut	-	grams of urea, 111 grams of rock phosphate and 117 grams of				
		muriate of potash per arecanut plant.				
Animal Husbandry: Precautions to be taken to avoid leptospirosis and ectoparasitic						

Animal Husbandry: - - Precautions to be taken to avoid leptospirosis and ectoparasitic infestations. Care should be taken to avoid fungal growth in cattle feed. Deworm the animals after dung examination. (CAADECCS, KVASU, Mannuthy).