



## **Agromet Advisory Bulletin for Thrissur District**





Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 47/2023, (94) Friday

**Dated 24.11.2023** 

	Observed 8 <sup>th</sup> Nove		U	st week ovember 2	2023)		Weather Forecast Valid for 25 <sup>th</sup> November to 29 <sup>th</sup> November 2023)						
18/11	19/11	20/11	21/11	22/11	23/11	24/11	Date		25/11	26/11	27/11	28/11	29/11
2.0	0.0	0.0	0.0	11.4	62.0	0.0	0.0Rainfall (mm)		10	5	3	3	2
34.1	33.8	33.9	33.0	34.6	33.1	32.5	T <sub>max</sub> ( <sup>0</sup> C)		33	33	33	33	33
24.5	25.2	23.1	24.0	24.8	24.2	25.9	$T_{min}(^{0}C)$		26	24	24	24	24
Partially Cloudy	Partially Cloudy	Cloudy	Partially Cloudy	Cloudy	Overcast	Cloudy	AM	Cloud cover	Overcast	Overcast	Overcast	Cloudy	Cloudy
Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy		PM						
95	77	81	80	95	98	77	RH-I (%)		96		96	96	96
55	55	55	57	64	67		RH-II (%)		71	71	71	71	71
1.8	1.9	2.5	1.5	1.9	1.6	2.3	Wind Sp	peed (km/hr.)	2	2	4	2	2
C	ENE	C	C	ENE	C	E	AM	Wind direction	SW	wsw	NW	W	W
SSE	NE	NW	ENE	NW	NNE		PM						
Rainfall (mm) in last week				Rain	Rainfall (mm) from 01/01/2023 to till dated					Rainfall (mm) from 01/10/2023 to today			
75.4				2084.4					465.4				

Weather summary / Alert: As per the forecast of IMD, there is a chance of moderate rainfall in isolated places of the district. Above normal rainfall is expected in Kerala.

## C. General and Agrometeorological Advisories for Thrissur District

Prophylactic measures can be taken to control and to avoid the spread of crop pest and diseases. Proper mulching can be followed to conserve soil moisture.

Crop	Pest/ Disease	Crop/ Animal Advisory
Rice (Mundakan) Tillering stage	Bacterial Leaf blight	To control and prevent bacterial leaf blight in paddy, dissolve 20 g of cow dung in one litre of water and spray the supernatant liquid by mixing with 20g of pseudomonas.
Vegetables (chilly)	Anthracnose disease	There is a chance of Anthracnose disease in chilly. As a prophylactic measure, apply 1% Bordeaux mixture or spray pseudomonas 20 gm per liter.
Turmeric	Leaf Blotch	Due to high relative humidity, there is a chance of leaf blotch diseases in turmeric. As a prophylactic messure, spray 2ml hexaconazole or 1 ml Propiconazole or 2 gram SAAF or 2.5 g copper oxychloride or 2 g copper hydroxide dissolved in one litre of water. Otherwise 1% Bordeaux mixture can be sprayed. Spraying can be continued at fortnightly intervals depending upon the intensity of the disease. Uproot and destroy fully infected plants.

Animal Husbandry: Vaccinate animals against foot and mouth disease and hemorrhagic septicemia. If needed give medicine agaist worm infestation after dung examination. (CAADECCS, KVASU, Mannuthy).