



## Gramin Krishi Mousam Sewa (GKMS), Thrissur



## Agromet Advisory Bulletin for Thrissur District (Valid from 08.12.2021 to 12.12.2021)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 50/2021, (99) Tuesday

dated 07.12.2021

A. Significant weather for the preceding week at vellanikkara							
Rainfall	Maximum	Minimum	RH	RH	Wind speed	Sun shine	Evap. (mm)
(mm)	Temp.(°C)	Temp.(°C)	FN (%)	AN (%)	(Km/h)	hours	
(20)	(31.3-33.4)	(22.8-25.2)	(79-96)	(54-71)	(1.0-7.0)	(1.4-9.6)	(2.3-3.5)

B. Medium range weather forecast of IMD for next five days					
Weather parameters	08.12.2021	09.12.2021	10.12.2021	11.12.2021	12.12.2021
Rainfall (mm)	5	4	2	1	1
Maximum temperature (°C)	33	33	33	33	33
Minimum temperature (°C)	23	23	23	23	23
Maximum Relative Humidity (%)	89	89	89	89	89
Minimum Relative Humidity (%)	70	70	70	70	70
Wind speed (kmph)	8	8	10	10	8
Wind direction (deg)	90	50	70	50	50
Total cloud cover (octa)	3	4	3	4	5

Weather summary / Alert: light rainfall is expected in the district.

Above normal rainfall is expected in Kerala.

RAINFALL FORECAST FOR NEXT FIVE DAYS					
<b>2021 December 08</b>	2021 December 09	2021 December 10	2021 December 11	2021 December 12	
Light rain	Light rain	Light rain	Light rain	Light rain	

## C. General and Agrometeorological Advisories for Thrissur District

The wind speed may increase. Give strong propping to banana and staking to vegetables.

Crop	Pest/ Disease		Crop/ Animal Advisory		
Mundakan Rice in Panicle initiation	Brown Plant Hopper (BPH)		During cloudy weather condition, there is a chance of Brown plant hopper attack in rice. Spray Imidacloprid 3ml per ten litre of water or Thiamethoxam 2 gram per ten litres of water.		
Coconut	Bud rot		To control the Bud rot in coconut, chisel out the affected area and apply10% Bordeaux paste. Care should be taken to protect the affected area from heavy rain.		
Turmeric	Leaf Blotch (Taphrina maculans)		Due to high relative humidity, there is a chance of leaf blotch diseases in turmeric. Remove and destroying disease affected plants. As a prophylactic measure, spray 2ml hexaconazole (Contaf) or 2 gram SAAF dissolved in one litre of water. Spraying can be continued at fortnightly intervals depending upon the intensity of the disease.		
Animal	Deworm the animals after dung examination. Vaccinate animals against foot and mouth disease.				
Husbandry	Care should be taken to avoid ecto parasitic infestation. (CAADECCS, KVASU, Mannuthy).				