



**Kerala Agricultural University, Vellanikkara, Thrissur**  
**Department of Agricultural Meteorology, College of Horticulture**

**Email: cohagmet@kau.in & kauagmet@yahoo.co.in**

**Block Level Experimental Agromet Advisory Bulletin**

**(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)**



**WADAKKANCHERY BLOCK, THRISSUR DISTRICT**

**AAS 41/2018(82), Friday**

**12.10.2018**

**The Agricultural advisory of this bulletin is not applicable in flood-affected areas.**

**SIGNIFICANT WEATHER FOR THE PRECEDING WEEK AT VELLANIKKARA:**

The total rainfall received 106.8 mm. the maximum temperature ranges from 32.0°C to 34.0°C, minimum temperature ranges from 22.6°C to 25.2°C. Morning relative humidity ranges from 80% to 96% and afternoon relative humidity ranges from 63% to 81%. Wind speed ranges from 0.5 to 7.5 km/h. Bright sunshine hours range from 0.4 to 8.9 hours. Evaporation ranges from 1.8 to 3.7 mm.

**CURRENT SYNOPTIC SITUATION:** A Deep Depression lies over Odisha centred near Lat 20.6° N/Long. 84.9° E. A severe cyclonic storm (LUBAN) lies over west Central Arabian sea near Lat 14.7° N/Long. 57.4° E. A trough runs from the deep depression to Kerala across Coastal Andhra Pradesh & Tamil Nadu and extends upto 3.1km Above mean sea level.

**WEATHER FORECAST FOR THE NEXT FIVE DAYS:**

Weather parameters	13.10.2018	14.10.2018	15.10.2018	16.10.2018	17.10.2018
Rainfall (mm)	4	4	10	0	14
Maximum temperature (°C)	33.4	33.2	33.4	33.2	33.0
Minimum temperature (°C)	24.6	24.2	24.4	24.6	24.0
Total cloud cover (octa)	6	6	6	6	6
Maximum Relative Humidity (%)	90	90	90	90	90
Minimum Relative Humidity (%)	70	70	70	70	70
Wind speed (kmph)	006	006	006	006	006
Wind direction (deg)	250	250	250	250	250

**As per IMD forecast, light rainfall in isolated places is expected for coming days in Wadakkanchery Block**

**AGRO-METEOROLOGICAL ADVISORIES:**

Crop	Crop stage	Pest/ Disease	Advisories
Paddy	Harvesting stage	-	First paddy crop is in Harvesting stage. Land preparation for second paddy crop can be started.
Coconut	-	Rhinoceros beetle	After thorough cleaning of the crown region, apply cartap hydrochloride 4G @ 20 g mixed with fine sand 200 g to the inner most leaf axils surrounding the spindle leaf or a mixture of fipronil 0.3 G @ 20 g and fine sand @ 200 g to the inner most leaf axils. This application has to be repeated thrice a year. (January, May and September/ October) for better results.
Banana	-	Spodoptera	Destroy the affected leaves along with the spodoptera. For severe attack spray quinalphos 2ml per one litre of water or chlorantraniliprole 3 ml/10 litre of water.
Animal Husbandry	<ul style="list-style-type: none"> <li>Animal should not be left to graze under hot sun from morning 10.00 AM to evening 3.00 PM.</li> <li>Concentrate feed should be sun dried regularly to avoid mould growth</li> <li>Deworm the animals and birds and vaccinate them against infectious diseases. (CAADECCS, KVASU, Mannuthy)</li> </ul>		